


ARMY INST OF DENTAL RESEARCH WASHINGTON D C
PERIODONTOSIS: A CASE REPORT WITH SCANNING ELECTRON MICROSCOPIC--ETC(U) AD-A036 133 FEB 77 J M BRADY, A L ALLEN UNCLASSIFIED NL LOF ! AD A036133 END 110 DATE 3-77





Periodontosis: A Case Report With Scanning Electron Microscopic Observations




John M. Brady, B.S., D.D.S., M.S.P.H.+

Andrew L. Allen, B.A., D.M.D., M.S.*

COPY AVAILABLE TO DDC DOES NOT PERMIT FULLY LEGIBLE PRODUCTION





+Colonel, D.C., Chief, Department of Biophysics, United States Army Institute of Dental Research, Walter Reed Army Medical Center, Washington, D.C. 20012

*Assistant Professor, Department of Periodontics, Baltimore College of Dental Surgery, Dental School, University of Maryland, 666 West Baltimore Street, Baltimore, Maryland 21201 Periodontosis: A Case Report with
Scanning Electron Microscopic Observations

Recent studies have suggested that rather specific bacterial populations may be related to varying periodontal disease states. 1-3 Periodontosis, as defined by Baer, 4 has been associated with deep periodontal defects which contain a high percentage of gram-negative anaerobic rods.3-5 Isolates of these rods have induced extensive alveolar bone loss accompanied by a marked osteoclastic response in germ-free rats.6,7 Rather precise sampling techniques3,8 and electron microscopic observations on extracted teeth have provided circumstantial evidence that the gram-negative organisms are located near the apex of the advancing lesion of periodontosis in humans. More information regarding the location and morphological characteristics of the subgingival organisms of periodontosis would be helpful in determining the possible cause and effect relationship between these bacteria and the clinical disease itself. The Scanning Electron Microscope (SEM) has been shown to be an excellent instrument for the examination of naturally occurring plaque structure, 9 yet no reports could be found concerning its use in evaluation of the periodontosis lesion. The purpose of this case report is to present the clinical finidngs and SEM observations of a tooth and associated periodontal tissues from a patient with a diagnosis of periodontosis.

Case Report

This 21-year old black female presented to Epes Army Dental Clinic for treatment complaining of "sore lower gums." Past medical history revealed an anemia of 5-6 years duration which has recently been diagnosed as hereditary elliptocytosis associated with minimal hemolysis. The patient recalled that spaces began developing between her front teeth at age 16 and previous dental extractions had occurred after the teeth had become "loose and sore."

Oral examination revealed normal mucosal surfaces and physiologic pigmentation of the gingival tissues. Teeth numbers 2, 3, and 30 were missing. Examination with the periodontal probe revealed severe vertical defects on numbers 9, 14, 24, 25, and 30. Plaque and calculus were evident, especially interproximally. Clinical photographs at the initial visit and an abbreviated radiographic survey may be seen in Figures 1 and 2. The patient's 19-year old sister presented similar clinical and radiographic findings.

Local factors, while present, did not appear commensurate with the amount of localized periodontal destruction in the molar and incisor regions. The treatment plan included the extraction of tooth numbers 9, 24, and 25 which were felt to have a hopeless prognosis. A periodontal charting of the maxillary incisors six weeks prior to extraction is shown in Table 1.

Materials and Methods

Tooth #9 was extracted securing a conservative intact V-shaped section of palatal tissue following the technique described by Ramfjord. 11 A soft tissue portion of the distal periodontal defect remained attached following extraction. The specimen was placed in chilled 2.5% glutaraldehyde in cacodylate buffer, pH 7.4, for 24 hours. After fixation the tooth was decalcified in EDTA for six weeks, at 40C, then sliced into several longitudinal sections through the intact attached soft tissue and root surface. The soft tissue was gently reflected from the lingual surface of the tooth the tooth so as to minimize distortion of the pocket contents and the attached soft tissues. The speicmens were washed in buffer, post-fixed in in assium tetroxide in buffer, dehydrated in increasing concentrations of alcohol, placed in any acetate and critical-point-dried in liquid carbon dioxide. The tooth with contiguous soft tissue was mounted on a sample stub, evaporation-coated with a thin layer of gold and palladium and examined in a scanning electron microscope.*

Results

A representative longitudinal section of tooth #9 with attached soft tissue was examined in the SEM (Figure 3). The soft tissue was observed to be attached to the lingual surface of decalcified root at the base of the periodontal pocket (Fig. 3).

^{*} Model AMR-1000 Scanning Electron Microscope. Advanced Metals Research Corporation; 160 Middlesex Turnpike; Bedford, MA 01730

The attachment of the fibrous soft tissue of the periodontal ligament was clearly evident in the photomicrographs (Fig. 4). An extensive examination of the entire dental surface of the periodontal defect was made with the SEM. The rock surface of the tooth was covered with scattered clumps of microorganisms of various shapes including rods, cocci and filaments. The apical portion of the dental surface of the periodontal defect was sparsely populated by red blood cells as observed at low magnification (Fig. 5). At higher magnifications rod-shaped microorganisms of uniform morphology were observed to be present on the cemental surface and on the disrupted soft tissues at the base of the defect (Figs. 6 and 7). These rod-shaped organisms were present in large clumps at the junction between the root surface and attached ligament fibers, and distributed evenly across the apicalmost 0.44 mm of the root surface of the periodontal defect. Examination of the entire 0.2 mm² area just coronal to the ligament attachment and of the adjoining soft tissue in the SEM did not reveal any microorganism morphology other than these short rods.

Discussion

Baer has defined periodontosis as a disease of the periodontium occurring in an otherwise healthy adolescent, which is characterized by a rapid loss of alveolar bone about more than one tooth of the permanent dentition where the periodontal destruction is not commensurate with the amount of local irritants present. The clinical characteristics and past dental history of this patient conform in most respects to this definition. The medical finding

of hereditary elliptocytosis was an interesting one. Hereditary elliptocytosis is transmitted as an autosomal dominant trait affecting the erythrocytes causing them to be elliptical or oval in shape. In mild forms, it is apparently a harmless disorder, although occassionally the disease is associated with severe hemolysis. 12 The mild elliptocytosis present in this patient was probably an incidental finding because a review of the literature failed to reveal any previous reports associating the two diseases.

Based upon the probable age of onset, the similar involvement of a sibling, and the clinical and radiographic findings, the diagnosis of advanced periodontosis was made. Disruption of the pocket contents undoubtedly occurred during reflection of the soft tissues from the root surface. This disruption, however, would be unlikely to cause displacement of organisms in a coronalapical direction. Therefore, the consistent finding of an almost pure culture of rod-shaped organisms in the apical portion of the defect, continuous with the soft tissue components is highly suggestive of their actual location within the pocket. It was not possible to determine whether the rod-shaped organisms seen in this study were identical with the organisms reported by Newman et al, 3 nor was it possible to distinguish morphologic groups of organisms. However, these findings specifically locate rods at the apex of the lesion and provides additional evidence that the lesion of periodontosis is characterized by a rather distinct microbial population. If this microbiologic component can be shown to be a major factor in the pathogenesis of this disease, perhaps both the "classical" molar-incisor involvement and the more generalized form of periodontosis

should be referred to as juvenile periodontitis. 13

Summary

A tooth and associated periodontal tissues from a patient with the diagnosis of periodontosis was subjected to scanning electron microscopic evaluation after reflection of the soft tissue portion of the lingual defect. The cemental surface of the apical portion of the lesion was found to be heavily populated by markedly similar rod-shaped organisms. The microorganisms were observed to be continuously present in a coronal-apical direction as the transition from cementum to the attached soft tissues at the base of the defect were examined. These findings provide additional evidence that the lesion of periodontosis is characterized by a rather distinct microbial population of rod-shaped organisms located predominately at the base of the defect.

REFERENCES

- 1. Listgarten, M.A.: Structue of the microbial flora associated with periodontal health and disease in man, a light and electron microscopic study. J Periodontol 47:1, 1976.
- Darwish, S.; Hyppa, T.; Socransky, S.S.: Predominate cultivatable microorganisms in periodontitis and periodontosis. II. Early periodontitis. J Dent Res 52:289, 1973.
- Newman, M.G.; Socransky, S.S.; Savitt, E.D.; Propas, D.A.; and Crawford, A.: Studies of the microbiology of periodontosis. J Periodontol 47:373, 1976.
- Baer, P.N.: The care for periodontosis as a clinical entity. J Periodontol 42:516, 1971.
- 5. Newman, M.G.; Socransky, S.S.; and Listgarten, M.A.: Relationship of microorganisms to the etiology of periodontosis. <u>J</u> <u>Dent Res</u> 53:324, 1974.
- 6. Irving, J.I.; Newman, M.G.; Socransky, S.S.; and Heeley, J.D.: Histologic changes in experimental periodontal disease in rats mono-infected with a gram-negative organism. Arch Oral Biol 20:219, 1975.
- 7. Irving, J.T.; Socransky, S.S.; Newman, M.G.; and Savitt, E.D.: Periodontal destruction induced by <u>Capnocytophaga</u> in gnotobiotic rats. J Dent Res 55:B783, 1976.
- 8. Crawford, A.; Darwish, S.; Newman, M.; Manganiello, A.D.; and Socransky, S.S.: Predominant cultivable microorganisms in periodontitis and periodontosis I. Sampling, cultivation, characterization. J Dent Res 52:288, 1973.
- 9. Newman, H.N.: Structure of approximal human dental plaque as observed by Scanning Electron Microscopy. Arch Oral Biol 17:1445-1453, 1972.
- Fletcher, Eugene C., MAJ, United States Army Medical Corps. Personal correspondence. 1975.
- Ramfjord, S. and Costich, E.R.: Healing after simple gingivectomy. <u>J Periodont</u> 34:401, 1963.
- 12. Harrison's Principles of Internal Medicine, McGraw-Hill Book Company, New York, N.Y. Chapter 338, p.1615, Sixth Edition, 1971.
- 13. Fourel, J.: Periodontosis, juvenile periodontitis, or Gottlieb Syndrome: Report of four cases. <u>J Periodont</u> 45:234, 1974.


TABLE

Results of periodontal probing six weeks prior to extraction of tooth number 9 and examination of tissue (pocket depth in mm.).


FIGURES

- FIGURE 1: Photographs of maxillary and mandibular incisor region showing migration and soft tissue destruction.
- FIGURE 2: Periapical radiograph demonstrating loss of osseous support in the incisor and molar areas. Previous extractions were for periodontal reasons.
- FIGURE 3: Scanning Electron Micrograph (SEM) of the extracted, decalcified root segment with attached soft tissue. The soft tissue wall of the periodontal defect (a) has been reflected away from the cemental surface (b). Magnification 24 times.
- FIGURE 4: SEM of the apical region of the periodontal defect.


 The base of the defect is delineated by the presence of the periodontal ligament (arrow). Magnification 118 times.
- FIGURE 5: SEM of the apical portion of the cemental surface of the periodontal defect. Occasional red blood cells are on the surface. The fibrous base of the periodontal ligament is marked by the arrow. Magnification 235 times.
- FIGURE 6: SEM of the cemental surface at the base of the periodontal defect. Short rod-like organisms cover the cementum and occasional red blood cells. Magnification 6000 times.
- FIGURE 7: SEM of the coronal-most tissue at the base of the defect. Short rod-like organisms are adjacent to these tissues. Magnification 12,000 times.








	fa	ci	a l	lir	gu	al
#	E	f	d	m	f	d
7	2	2	2	2	. 1	2
8	3	1	2	3	3	2
•	6	2.	8	7	5	7
10	4	2	3	4	4	3





SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered) READ INSTRUCTIONS BEFORE COMPLETING FORM REPORT DOCUMENTATION PAGE . REPORT NUMBER 2. JOYT ACCESSION NO. 3. RECIPIENT'S CATALOG NUMBER E (and Subtitle) T & PERIOD COVERED Periodontosis: A Case Report with Scanning - Nov 1076 Electron Microscopic Observations . 8. CONTRACT OR GRANT NUMBER(*) JOHN M BRADY ANDREW L ALLEN PERFORMING ORGANIZATION NAME AND ADDRESS 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK/UNIT NUMBERS US Army Institute of Dental Research 672775.25.7.00622 1498# DAOH6038 Walter Reed Army Medical Center Washington, DC 20012 11. CONTROLLING OFFICE NAME AND ADDRESS REPORT DATE US Army Medical Research and Development Command-15 Feb 77 ATTN: (SGRD-RP) WASH DC 20314 4. MONITORING AGENCY NA ADDRESS(If different from Controlling Office) 15. SECURITY CLASS. (of this report) UNCLASSIFIED 15a. DECLASSIFICATION/DOWNGRADING 16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited. 17. DISTRIBUTION STATEMENT (of the obstract entered in Block 20, If different from Report) 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Periodontosis, scanning electron microscopy ABSTRACT (Continue on reverse side if necessary and identity by block number) canning microscopy was performed on the root surface of tooth removed due to severe periodontal destruction as a result of periodontosis. The cemental surface was covered with markedly similar rod shaped microorganisms. These findings provide additional evidence that this disease is caused by a characteristic bacteria, located at the base of the advancing lesion.

UNCLASSIFIED (BEISTIN STATE CHARLES OF THE ROLL OF THIS PAGE (

DD 144 72 1473 EDITION OF I NOV SE IS OBSOLETE

	NA DELASSIFICATION OF THE PAGE ROSE DATE HAVE DESCRIBED (Described along)
BEFORE COMPLETING FORM	REPORT DOGUMENTATION PAGE
S RECIPIONTE CATALOG RUMBER	REPORT HUMBER
DAMANOS COMPAS A PARANO CONTRACO	(withdays bon) advar
THE PROPERTY OF SHAPE TO SHAPE	Periodontosis: A Case Report with Scanning []
E CONTRACT OR GRANT NUMBERS	YOUR Y GRADE
	LATIVITY MACH
PERCHANGLENGER PROJECT TASK AREA WORK UNIT NUMBERS 0727/0.25.7.00022 1.988 JACOBO 33	Seronmun cadaquization name and abbases 15 Arroy Institute of Dental Rasearch talter Reed Army Medical Center tashinuton, UC 20012
AL REPORT DANG TO FAD 11 TO HOMOTER OF PLACES	CONTROLLING OFFICE NAME AND ADDRESS JS Army Medical Research and Development Command TTR: (JGRU-RF)
TO SECURITY CLASS (of Diffe report)	NON TO COUNTY NEW A NOUTESSAIL THE CONTESTING THE OF THE O
DHICH ASSET FLOW DOWNGRADING	10 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
	COSTRIBUTION BY ATEMENT (of this Papert)
	This document has been approved for public release is unlimited.
n Report)	DISTRIBUTION STATEMENT (2) the southered in Plack 20, it different from
	ESTON YPATHS JERUS .
	(NEY WORDS (Continue on reverse tide if necessary and identify by block number)
	Periodontosis, scanaing electron microscopy
nce of thath removed due to stantosis. The cemental of microproanisms. These hase is caused by a character	eniodoncosis, scanning electron microscopy AMSYRACT (Control on revolve side in processes and identity of block home) I canning microscopy was performed on the root surfacevery periodonical decrination as a result of period artace was covered with markedly similar rod subpointings provide additional evidence that this disciplination becteria, located at the base of the advanctor

#	facial			lingual		
	m	f	d	m	f	d
7	2	2	2	2	1	2
8	3	1	2	3	3	2
9	6	2	8	7	5	7
10	4	2	3	4	4	3